

## Residual activity of $\alpha$ -chymotrypsin in water-1,2-propanediol mixtures

Komissarov I., Kuchierskaya A., Sirotkin V.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

### Abstract

© 2017 Nova Science Publishers, Inc. All rights reserved. A novel experimental method was proposed to study the stability of enzymes in water-organic mixtures. This method is based on the analysis of the residual enzyme activity. Advantages of our method: (I) The residual activity curves can be determined in the entire range of the water content in organic liquids. (II) Enzyme activity values are measured at fixed reaction conditions. (III) Information on the changes in the state of the catalytically active site can be obtained separately from the other contributions (for example, changes in the solvation of the reagents and products in various water - organic mixtures). To show the efficiency of the proposed method we studied the residual activity of bovine pancreatic  $\alpha$ -chymotrypsin (CT) in water-1,2-propanediol mixtures in the entire range of water content at 25°C. The obtained results show that the degree of stabilization/destabilization of  $\alpha$ -chymotrypsin depends strongly on the water content in organic solvent. (1) At high water content, the residual activity values are higher than 100%. (2) At low water content, the residual catalytic activity is ~90-95%, compared with that observed after incubation in pure water. (3) A minimum on the residual activity curve was observed at intermediate water content.

---

### Keywords

1,2-propanediol (1,2-PG), Biocatalysis in water-organic mixtures, Bovine pancreatic  $\alpha$ -chymotrypsin (CT), Organic solvent, Residual activity, Stability, Water

### References

- [1] Klibanov, A. M., (2001) Improving enzymes by using them in organic solvents, *Nature* 409, 241-246.
- [2] Carrea, G., Riva, S., (2000) Properties and synthetic applications of enzymes in organic solvents, *Angew. Chem. Int. Ed.* 39, 2226-2254.
- [3] Halling, P. J., (2004) What can we learn by studying enzymes in nonaqueous media? *Phil. Trans. R. Soc. Lond. B Biol. Sci.* 359, 1287-1297.
- [4] Micaelo, N. M., Soares C. M., (2007) Modeling hydration mechanisms of enzymes in nonpolar and polar organic solvents, *FEBS J.* 274, 2424-2436.
- [5] Clark, D. S., (2004) Characteristics of nearly dry enzymes in organic solvents: implications for biocatalysis in the absence of water, *Phil. Trans. R. Soc. Lond. B Biol. Sci.* 359, 1299-1307.
- [6] Serdakowski, A. L., Dordick, J. S., (2007) Enzyme activation for organic solvents made easy, *Trends Biotechnol.* 26, 54-48.
- [7] Rariy, R., Klibanov, A. M., (1997) Correct protein folding in glycerol, *Proc. Natl. Acad. Sci. US* 94, 13520-13523.
- [8] Sirotkin, V. A., Hüttel, R., Wolf, G., (2008) Enzyme-catalysed hydrolysis of L-amino acid esters in a low water organic solvent studied by isothermal calorimetry, *J. Therm. Anal. Calorim.* 93, 515-520.

- [9] Gonzalez Martinez, S., Alvira, E., Vergara-Cordero, L., Ferrer, A., Montañes-Clemente, I., Barletta, G., (2002) High initial activity but low storage stability observed for several preparations of subtilisin Carlsberg suspended in organic solvents, *BiotechnolProg.* 18, 1462-1466.
- [10] Castillo, B., Pacheco, Y., Al-Azzam, W., Griebenow, K., Devi, M., Ferrer, A., Barletta, G., (2005) On the activity loss of hydrolases in organic solvents: I. Rapid loss of activity of a variety of enzymes and formulations in a range of organic solvents, *J. Mol. Catalysis B: Enz.* 35, 147-153.
- [11] Fernandez, J. F. A., Halling, P., (2002) Operational stability of high initial activity protease catalysts in organic solvents, *Biotechnol. Prog.* 18, 1455-1457.
- [12] Sirotkin, V. A., Hütt, I R., Wolf, G., (2005) Thermochemical investigations of hydrolysis of p-nitrophenyl acetate in water-acetonitrile mixtures, *Thermochim. Acta* 426, 1-6.
- [13] Ke, T., Klibanov, A. M., (1998) On enzymatic activity in organic solvents as a function of enzyme history, *Biotechnol. Bioeng.* 57, 746-750.
- [14] Partridge, J., Hutcheon, G. A., Moore, B. D., Halling P. J., (1996) Exploiting hydration hysteresis for high activity of cross-linked subtilisin crystals in acetonitrile, *J. Amer. Chem. Soc.* 118, 12873-12877.
- [15] Zaks, A., Klibanov, A. M., (1988) The effect of water on enzyme action in organic media, *J. Biol. Chem.* 263, 8017-8021.
- [16] Yamane, T., Kojima, Y., Ichiryu, T., Shimizu, S., (1988) Biocatalysis in a microaqueous organic solvent, *Ann. NY Acad. Sci.* 542, 282-293.
- [17] Lee, C. S., Ru, M. T., Haake, M., Dordick, J. S., Reimer, J. A., Clark, D. S., (1998) Multinuclear NMR study of enzyme hydration in an organic solvent, *Biotechnol. Bioeng.* 57, 686-693.
- [18] McMinn, J. H., Sowa, M. J., Charnick, S. B., Paulaitis, M. E., (1993). The hydration of proteins in nearly anhydrous organic solvent suspensions, *Biopolymers.* 33, 1213-1224.
- [19] Kijima, T., Yamamoto, S., Kise, H., (1996) Study of tryptophan fluorescence and catalytic activity of  $\alpha$ -chymotrypsin in aqueous organic media, *Enz. Microb. Technol.* 18, 2-6.
- [20] Khmelnitsky, Yu., Mozhaev, V. V., Belova, A. B., Sergeeva, M. V., Martinek, K., (1991) Denaturation capacity: a new quantitative criterion for selection of organic solvents as reaction media in biocatalysis, *Eur. J. Biochem.* 198, 31-41.
- [21] Simon, L. M., Kotorman, M., Garab, G., Laczko, I., (2001) Structure and activity of b{cyrillic}-chymotrypsin and trypsin in aqueous organic media, *Biochem. Biophys. Res. Comm.* 280, 1367-1371.
- [22] Sirotkin, V. A., Mukhametzyanov, T. A., (2006) Effect of acetonitrile on the binding of competitive inhibitor proflavin and on the catalytic activity of bovine pancreatic  $\alpha$ -chymotrypsin, *Russ. J. Phys Chem. A.* 80, 803-808.
- [23] Sirotkin, V. A., Mukhametzyanov, T. A., Komissarov, I. A., (2009) Effect of tetrahydrofuran on the binding of the competitive inhibitor proflavin and the storage stability of bovine pancreatic  $\alpha$ -chymotrypsin, *Eng. Life Sci.* 9, 82-88.
- [24] Sirotkin, V. A., Mukhametzyanov, T. A., Karmanova, Y. V., (2007) Effect of dioxane on the binding of competitive inhibitor proflavin and catalytic activity of bovine pancreatic  $\alpha$ -chymotrypsin. *Russ. J. Phys Chem. A.* 81, 1160-1164.
- [25] Ramanathan, K., Jonsson, B. R., Danielsson, B., (2000) Thermometric sensing of peroxide in organic media. Application to monitor the stability RBP-Retinol-HRP complex, *Anal. Chem.* 72, 3443-3448.
- [26] Fatibello-Filho, O., Cruz Viera, I., (2000) Construction and analytical application of a biosensor based on stearic acid-graphite powder modified with sweet potato tissue in organic solvents, *Fresenius J. Anal. Chem.* 368, 338-343.
- [27] Chang, N., Klibanov, A. M., (1992) Protein chromatography in neat organic solvents. *Biotechnol. Bioeng.* 39, 575.
- [28] Chang N., Hen S. J., Klibanov A. M., (1991) Protein separation and purification in neat dimethyl sulfoxide. *Biochem. Biophys. Res. Comm.* 176, 1462.
- [29] Bromberg, L. E., Klibanov, A. M., (1995) Transport of proteins dissolved in organic solvents across biomimetic membranes. *Proc. Natl. Acad. Sci. USA* 92, 1262.
- [30] Gregory, R. B. (1995) Protein hydration and glass transition behavior. In: *Protein-Solvent Interactions* (Gregory, R. B., ed.), Marcel Dekker, New York, 191-264.
- [31] Rupley, J. A., Careri, G., (1991) Protein hydration and function. *Adv. Protein Chem.* 41, 37-172.
- [32] Kuntz, I. D., Kauzmann, W., (1974) Hydration of proteins and polypeptides. *Adv. Protein Chem.* 28, 239-345.
- [33] Bull, H. B. (1944) Adsorption of water vapor by proteins. *J. Amer. Chem. Soc.* 66, 1499-1507.
- [34] Luscher-Mattli, M., Ruegg, M., (1982) Thermodynamic functions of biopolymer hydration. I. Their determination by vapor pressure studies, discussed in an analysis of the primary hydration process. *Biopolymers* 21, 403-418.
- [35] Luscher-Mattli, M., Ruegg, M., (1982) Thermodynamic functions of biopolymer hydration. II. Enthalpy-entropy compensation in hydrophilic hydration process. *Biopolymers* 21, 419-429.

- [36] Bone, S. (1987) Time-domain reflectometry studies of water binding and structural flexibility in chymotrypsin. *Biochem. Biophys. Acta.* 916, 128-134.
- [37] Hnojewy, W. S., Reyerson, L. H., (1961). Further studies on the sorption of HO and DO vapors by lysozyme and the deuteriumhydrogen exchange effect. *J. Phys. Chem.* 65, 1694-1698.
- [38] Sirotkin V. A., Korolev D. V., (2005) Effect of acetonitrile on the hydration of human serum albumin films: a calorimetric and spectroscopic study. *Thermochim. Acta*, 432, 246-253.
- [39] Sirotkin, V. A., Khadiullina, A. V., (2011) Hydration of proteins: excess partial enthalpies of water and proteins. *J. Phys. Chem. B*, 115, 15110-15118.
- [40] Sirotkin, V. A., Komissarov, I. A., Khadiullina, A. V., (2012) Hydration of proteins: excess partial volumes of water and proteins, *J. Phys. Chem. B*, 116, 4098-4105.
- [41] Sirotkin, V. A., Khadiullina, A. V., (2014) A study of the hydration of ribonuclease A using densitometry: Effect of the protein hydrophobicity and polarity. *Chem. Phys. Lett*, 603, 13-17.
- [42] Sirotkin, V. A., Khadiullina, A. V., (2013) Gibbs energies, enthalpies, and entropies of water and lysozyme at the inner edge of excess hydration. *J. Chem. Phys*, 139, 075102/1-075102/9.
- [43] Sirotkin, V. A. (2005) Effect of dioxane on the structure and hydration/dehydration of  $\alpha$ -chymotrypsin as measured by FTIR spectroscopy. *Biochim. Biophys. Acta.*, 1750, 17-29.
- [44] Kocherbitov, V., Arnebrant, T., (2010) Hydration of Lysozyme: The Protein-Protein Interface and the Enthalpy-Entropy Compensation. *Langmuir*, 26, 3918-3922.
- [45] Foss, J. G., Reyerson, L. H., (1958) The sorption of water vapor by lyophilized ribonuclease. *J. Phys. Chem.* 62, 1214-1216.
- [46] Amberg C. H. (1957) Heats of adsorption of water vapor on bovine serum albumin. *J. Amer. Chem. Soc*, 79, 3980-3984.
- [47] Almog R., Schrier E. E., (1978). The dependence of the heat of solution of lyophilized solid ribonuclease A on water content. *J. Phys. Chem*, 82, 1701-1702.
- [48] Smith A. L., Shirazi H. M., Mulligan S. R., (2002) Water sorption isotherms and enthalpies by lysozyme using the quartz crystal microbalance/heat conduction calorimeter. *Biochim. Biophys. Acta*, 1594, 150-159.
- [49] Kocherbitov V., Arnebrant T., Söderman O., (2004) Lysozyme-water interactions studied by sorption calorimetry. *J. Phys. Chem. B*, 108, 19036-19042.
- [50] Bychkova, V. E., Dujsekina, A. E., Klenin, S. I., Tiktopulo, E. I., Uversky, V. N., Ptitsyn O. B., (1996) Molten globule-like state of s{cyrillic}ytochrome s{cyrillic} under conditions simulating those near the membrane surface, *Biochemistry* 35, 6058-6063.
- [51] Timasheff, S. N. (1992) Water as Ligand: Preferential binding and exclusion of denaturants in protein unfolding. *Biochem.* 31, 9857-9864.
- [52] Timasheff, S. N. (2002) Protein hydration, thermodynamic binding, and preferential hydration. *Biochem.* 41, 13473-13482.
- [53] Banipal T. S., Singh G., (2004) Thermodynamic study of solvation of some amino acids, diglycine and lysozyme in aqueous and mixed aqueous solutions. *Thermochim. Acta.* 412, 63-83.
- [54] Kumar, A., Attri, P., Venkatesu, P., (2012) Effects of polyols on the native structure of  $\alpha$ -chymotrypsin: A comparative study. *Thermochim. Acta.* 536, 55-62.
- [55] Kumar, A., Venkatesu, P., (2012) Overview of the stability of  $\alpha$ -chymotrypsin in different solvent media. *Chem. Rev.* 112, 4283-4307.
- [56] Gekko K., Timasheff S. N., (1981) Mechanism of protein stabilization by glycerol: Preferential hydration in glycerol-water mixtures. *Biochem* 20, 4667-4676.
- [57] Fersht A. Structure and mechanism in protein science: a guide to enzyme catalysis and protein folding. New York: Freeman & Co, 1999.
- [58] Lehninger A. L., Nelson D. L., Cox M. M. Principles of Biochemistry. New York: Worth, 1993.
- [59] Perrin, D. D., Armarego, W. L. F., Perrin, D. R., Purification of Laboratory Chemicals, Oxford: Pergamon Press, 1980.
- [60] Borisover, M. D., Sirotkin, V. A., Solomonov, B. N., (1993) Calorimetric study of human serum albumin in the water - dioxane mixtures. *J. Phys. Org. Chem.* 6, 251-253.
- [61] M. R. Eftink, R. E. Johnson, and R. L. Biltonen, The application of flow microcalorimetry to the study of enzyme kinetics, *Anal. Biochem.* 111 (1981) 305-320.
- [62] Atkins, P. W. (2006) Physical Chemistry. 8ed. Oxford: Oxford University Press.
- [63] Prausnitz, J. M. (1969) Molecular Thermodynamics of Fluid-Phase Equilibria. N. J.: Prentice-Hall, Inc., Engelwood Cliffs.
- [64] Belousov, V. P., Panov, M.Y. (1994) Thermodynamic properties of aqueous solutions of organic substances. Boca Raton, Fla.: CRC Press.

- [65] Koga, Y., Siu, W. W. Y., Wong, T. Y. H., (1990) Excess partial molar free energies and entropies in aqueous tert-butanol solutions at 25C. J. Phys. Chem. 94, 7700 - 7706.
- [66] Franks F., (1982) Hydrophobic interactions - a historical perspective, Faraday Symp. Chem. Soc., 17, 7-10.